



Year: 2019

Whole genome sequencing for drug resistance profile prediction in *Mycobacterium tuberculosis*

Gygli, Sebastian M ; Keller, Peter M ; Ballif, Marie ; Blöchliger, Nicolas ; Hömke, Rico ; Reinhard, Miriam ; Loiseau, Chloé ; Ritter, Claudia ; Sander, Peter ; Borrell, Sonia ; Loo, Jimena Collantes ; Avihingsanon, Anchalee ; Gnokoro, Joachim ; Yotebieng, Marcel ; Egger, Matthias ; Gagneux, Sebastien ; Böttger, Erik C

Abstract: Whole genome sequencing allows rapid detection of drug-resistant isolates. However, the availability of high-quality data linking quantitative phenotypic drug susceptibility testing (DST) and genomic data has thus far been limited. We determined drug resistance profiles of 176 genetically diverse clinical isolates from Democratic Republic of the Congo, Ivory Coast, Peru, Thailand and Switzerland by quantitative phenotypic DST for 11 antituberculous drugs using the BD BACTEC MGIT 960 system and 7H10 agar dilution to generate a cross-validated phenotypic DST readout. We compared DST results with predicted drug resistance profiles inferred by whole genome sequencing. Classification of strains by the two phenotypic DST methods into resistotype/wild type populations was concordant in 73-99 % of cases, depending on the drug. Our data suggests that the established critical concentration (5 mg/L) for ethambutol resistance (MGIT 960 system) is too high and may misclassify strains as susceptible, compared to 7H10 agar dilution. Increased minimal inhibitory concentrations were explained by mutations identified by whole genome sequencing. Using whole genome sequences, we were able to predict quantitative drug resistance levels for the majority of drug resistance mutations. Predicting quantitative levels of drug resistance by whole genome sequencing was partially limited due to incompletely understood drug resistance mechanisms. The overall sensitivity and specificity of whole genome-based DST were 86.8 % and 94.5 %, respectively. Despite some limitations, whole genome sequencing has the potential to infer resistance profiles without the need for time-consuming phenotypic methods.

DOI: <https://doi.org/10.1128/AAC.02175-18>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-168536>

Journal Article

Accepted Version



The following work is licensed under a Creative Commons: Attribution 4.0 International (CC BY 4.0) License.

Originally published at:

Gygli, Sebastian M; Keller, Peter M; Ballif, Marie; Blöchliger, Nicolas; Hömke, Rico; Reinhard, Miriam; Loiseau, Chloé; Ritter, Claudia; Sander, Peter; Borrell, Sonia; Loo, Jimena Collantes; Avihingsanon,

Anchalee; Gnokoro, Joachim; Yotebieng, Marcel; Egger, Matthias; Gagneux, Sebastien; Böttger, Erik C (2019). Whole genome sequencing for drug resistance profile prediction in Mycobacterium tuberculosis. Antimicrobial Agents and Chemotherapy, 63(4):e02175-18.
DOI: <https://doi.org/10.1128/AAC.02175-18>

1 **Title**2 **Whole genome sequencing for drug resistance profile prediction in**
3 ***Mycobacterium tuberculosis***4 **Authors**

5 Sebastian M. Gygli^{1,2,#}, Peter M. Keller^{12,#}, Marie Ballif⁵, Nicolas Blöchliger³, Rico
6 Hömke^{3,4}, Miriam Reinhard^{1,2}, Chloé Loiseau^{1,2}, Claudia Ritter^{3,4}, Peter Sander^{3,4}, Sonia
7 Borrell^{1,2}, Jimena Collantes Loo¹¹, Anchalee Avihingsanon^{6,7}, Joachim Gnokoro⁸, Marcel
8 Yotebieng⁹, Matthias Egger^{5,10*}, Sebastien Gagneux^{1,2*§}, Erik C. Böttger^{3,4*§}

9 [#]equal contribution as first authors; ^{*}equal contribution as last authors; [§]corresponding
10 authors

11 **Affiliations**

12 ¹Swiss Tropical and Public Health Institute, Basel, Switzerland

13 ²University of Basel, Basel, Switzerland

14 ³Institute of Medical Microbiology, University of Zürich, Zürich, Switzerland

15 ⁴National Center for Mycobacteria, University of Zürich, Zürich, Switzerland

16 ⁵Institute of Social and Preventive Medicine, University of Bern, Bern, Switzerland

17 ⁶HIV-NAT/Thai Red Cross AIDS Research Centre, Bangkok, Thailand

18 ⁷TB Research Unit, Department of Medicine, Faculty of Medicine, Chulalongkorn
19 University, Bangkok, Thailand

20 ⁸Centre de Prise en Charge de Recherche et de Formation, Yopougon, Abidjan, Côte d'Ivoire

21 ⁹College of Public Health, Ohio State University, Columbus, Ohio, USA

22 ¹⁰Centre for Infectious Disease Epidemiology and Research, University of Cape Town, South
23 Africa

24 ¹¹Instituto de Medicina Tropical "Alexander von Humboldt", Universidad Peruana Cayetano
25 Heredia, Lima, Peru

26 ¹²Institute for Infectious diseases, University of Bern, Bern, Switzerland

27

28 Corresponding authors

29 Prof. Erik C. Böttger, MD, University of Zürich, Institute of Medical Microbiology, National
30 Center for Mycobacteria, Gloriastrasse 28/30, 8006 Zürich, Switzerland; Phone: +41 44 634
31 2660; Fax: +41 44 634 49 06; email: boettger@imm.uzh.ch

32
33 Prof. Sebastien Gagneux, PhD, Swiss Tropical and Public Health Institute, Socinstrasse 57,
34 4002 Basel, Switzerland; Phone: +41 61 284 8369; Fax: +41 61 284 8101; email:
35 sebastien.gagneux@swisstph.ch

37 Key words

38 *Mycobacterium tuberculosis*, quantitative phenotypic drug susceptibility testing, whole
39 genome sequencing, drug resistance, drug resistance level prediction

40 Abstract

41 Whole genome sequencing allows rapid detection of drug-resistant *Mycobacterium*
42 *tuberculosis* isolates. However, the availability of high-quality data linking quantitative
43 phenotypic drug susceptibility testing (DST) and genomic data has thus far been limited.

44 We determined drug resistance profiles of 176 genetically diverse clinical *M. tuberculosis*
45 isolates from Democratic Republic of the Congo, Ivory Coast, Peru, Thailand and
46 Switzerland by quantitative phenotypic DST for 11 antituberculous drugs using the BD
47 BACTEC MGIT 960 system and 7H10 agar dilution to generate a cross-validated phenotypic
48 DST readout. We compared DST results with predicted drug resistance profiles inferred by
49 whole genome sequencing.

50 Classification of strains by the two phenotypic DST methods into resistotype/wild type
51 populations was concordant in 73-99 % of cases, depending on the drug. Our data suggests
52 that the established critical concentration (5 mg/L) for ethambutol resistance (MGIT 960
53 system) is too high and may misclassify strains as susceptible, compared to 7H10 agar
54 dilution. Increased minimal inhibitory concentrations were explained by mutations identified
55 by whole genome sequencing. Using whole genome sequences, we were able to predict
56 quantitative drug resistance levels for the majority of drug resistance mutations. Predicting
57 quantitative levels of drug resistance by whole genome sequencing was partially limited due

58 to incompletely understood drug resistance mechanisms. The overall sensitivity and
59 specificity of whole genome-based DST were 86.8 % and 94.5 %, respectively.

60 Despite some limitations, whole genome sequencing has the potential to infer resistance
61 profiles without the need for time-consuming phenotypic methods.

62 **Introduction**

63 Timely and accurate drug susceptibility testing (DST) of *M. tuberculosis* isolates is vital to
64 prevent the transmission of multidrug-resistant strains (MDR – resistance to rifampicin and
65 isoniazid) (1). The slow growth and stringent biosafety requirements of *M. tuberculosis* make
66 obtaining a full DST profile by culture-based techniques a matter of weeks or months. In
67 addition, culture-based DST is notoriously challenging for several drugs, e.g. pyrazinamide
68 and ethionamide due to poor drug solubility in commonly used culture media (2).

69 Drug resistance in *M. tuberculosis* is mainly conferred by chromosomal mutations in a few
70 genes (3), making it possible to detect drug resistance by sequencing these genes or probing
71 them by molecular hybridisation (4). Several commercial tests for the detection of resistance-
72 associated mutations are available, e.g. the GenoType MTBDRplus V2 (Hain Lifescience
73 GmbH, Nehren, DE) (5) and the AID TB Resistance Line Probe Assay (AID GmbH,
74 Strassberg, DE) (6). The World Health Organisation (WHO) endorses line probe assays and
75 the Xpert® MTB/RIF assay (Cepheid, Sunnyvale, CA, USA) for the detection of rifampicin
76 resistance as a surrogate marker for multidrug-resistance (7, 8). These molecular tests have
77 high sensitivities for drugs with established target(s) of resistance and for which only a few
78 mutations are responsible for most resistance *in clinico* (e.g. rifampicin, isoniazid) (4).
79 However, these molecular tests show low sensitivity for heteroresistant strains (concomitant
80 presence of wild type (wt) and mutant or multiple different resistant variants in patient
81 isolates), when frequencies of mutant variants drop below 5-50 % (9, 10). Furthermore, there
82 are no commercially available rapid tests for many drugs currently/prospectively in use (e.g.
83 bedaquiline, delamanid, linezolid, p-aminosalicylic acid) and the WHO only recently defined
84 *ad interim* critical concentrations for bedaquiline and delamanid for use with the BACTEC
85 MGIT 960 system (11, 12).

86 A wealth of genomic data on drug-resistant *M. tuberculosis* has become available in the past
87 years (13, 14). Unfortunately, quantitative phenotypic DST data are lacking for most of the
88 genetic data sets, necessary to infer phenotypes from genotypes. In addition, DST data are

89 often limited as the strains were classified as susceptible or resistant using at the WHO-
90 defined critical concentration (15). There is an urgent need to link genotypic and phenotypic
91 drug resistance readouts to obtain a better understanding of the mechanisms influencing the
92 evolution and spread of drug resistance in *M. tuberculosis* (3, 16).

93 WGS of clinical isolates allows for accurate identification of established chromosomal
94 mutations increasing the minimal inhibitory concentration (MIC) (13, 17, 18) and may
95 ensure adequate treatment in days instead of months. We compared whole genome-based
96 drug resistance profiles with two culture-based quantitative DST methods for a total of 11
97 drugs, including rifampicin, rifabutin, isoniazid, all WHO group B (streptomycin, kanamycin
98 A, amikacin and capreomycin), as well as selected group A (moxifloxacin), group C
99 (ethionamide) and group D (ethambutol, pyrazinamide) drugs (11).

100 **Material and methods**

101 ***M. tuberculosis* isolates**

102 The initial data-set consisted of 189 *M. tuberculosis* isolates. A subset of 61 strains was used
103 to establish the phenotypic DST methodology. These 61 strains were collected by the Swiss
104 National Center for Mycobacteria between 2004-2015, and represent a broad spectrum in
105 geographic origin and drug resistance profiles (19–21). We then applied the quantitative DST
106 methodology to 125 prospectively collected clinical isolates from clinics participating in the
107 International epidemiology Databases to Evaluate AIDS (IeDEA) (22) in Peru, Thailand,
108 Ivory Coast and the Democratic Republic of the Congo (supplementary [Table S3](#)). Thirteen
109 strains had to be excluded due to failed WGS (n = 4, failed library preparation due to low
110 DNA quality), irreproducible DST results (n = 1), no growth in the 7H10 agar dilution assay
111 (n = 3), duplication (n = 1), mixed cultures (n = 2, cross-contamination or patient infected
112 with multiple strains) or transmission clusters (n = 2). The final set consisted of 176 strains.

113 **Phenotypic DST**

114 MGIT 960- and 7H10 agar dilution-based phenotypic DST were performed as described
115 previously (19). Critical concentrations used for the classification of strains into
116 resistant/susceptible aim to predict clinical outcome, i.e. treatment failure if a given strain is
117 resistant at the critical concentration. However, critical concentrations should ideally be
118 defined on the basis of the epidemiological cut-off (ECOFF: The highest wt MIC observed in

119 absence of any detectable resistance mechanism (23)), treatment outcomes and pharmako-
120 kinetic and -dynamic data. As *M. tuberculosis* infections are treated with combination
121 therapy, outcome data for single drugs are difficult to obtain (24). This calls for definition of
122 critical concentrations solely based on the ECOFF (11). We therefore classified strains as
123 belonging to the resistotype/wt populations on the basis the detection of growth/no growth at
124 the ECOFF derived from our data (25). Table 1 lists the ECOFFs used, supplementary Table
125 S2 the drug concentrations tested with the MGIT 960 and 7H10 agar-dilution assays and
126 Table 2 the genes screened for mutations with WGS. Further details on how the phenotypic
127 DST assays were performed are available in the supplementary materials.

128 **Data analysis**

129 The categorical agreement between classification of strains into resistotype/wt populations
130 using MGIT 960 and 7H10 agar dilution was based on detectable growth at the ECOFF
131 (Table 1). The numerical variation between the two methods was quantified as the geometric
132 standard deviation (SD, given with its standard error) of the ratio MIC MGIT 960/MIC agar
133 dilution, expressed as a number of 2-fold dilutions and denoted by σ . The geometric SD was
134 computed by fitting a log-normal distribution to the ratio MIC MGIT 960/MIC agar dilution
135 as implemented in the R package fitdistrplus (v.1.0-9) (26). If the data was compatible with σ
136 = 0, the geometric standard deviation could not be estimated and was defined as “not
137 applicable” (NA). The approach is a generalization of the Bland and Altman method (27),
138 taking censoring of the data into account. Strains for which the MGIT 960 MIC and 7H10
139 agar dilution MIC were both left-censored or both right-censored were excluded since no
140 information on the ratio could be derived.

141 Goodman and Kruskal’s gamma was used to quantify the rank correlation between the two
142 methods. No correlation could be calculated if the variance for either method was 0 and
143 denoted with “not applicable” (NA).

144 Distributions of wt and mutant MICs were analysed qualitatively based on the results of
145 7H10 agar dilution. We divided the dataset into two groups: drugs for which the MIC
146 distributions of wt and mutant strains did not overlap, and those for which MIC distributions
147 overlapped.

148 Sensitivities and specificities of WGS-based resistance profile inference were calculated
149 based on the 7H10 agar dilution results for all drugs, except pyrazinamide– for which the

150 MGIT 960 results were used, based on growth/no growth at the ECOFF, derived from our
151 data and the presence or absence of a putative resistance-associated mutation.

152 **Defining cut-offs for high/low-level MICs**

153 The therapeutic window of a drug is defined as the concentration range within which a drug
154 is considered to be effective and safe to use (28). Mutations can increase the MIC beyond the
155 therapeutic window and render the drug clinically ineffective. Drugs may have large
156 therapeutic windows beyond the ECOFF. For these, MIC increases caused by mutations may
157 still be within the therapeutic window of a drug: these strains might still be treatable by
158 increasing the drug dose. We analysed the distribution of MICs of mutant strains, and
159 assessed if cut-offs for low-level (within the therapeutic window) and high-level (beyond the
160 therapeutic window) MICs were definable. There were sufficient data available to define
161 distinct cut-offs for low/high-level MICs for isoniazid, rifampicin, streptomycin and
162 amikacin. For mutations conferring resistance to other drugs assayed in this study, no distinct
163 separation into resistotype populations with high/low-level MICs was possible due to wide
164 ranges of MICs conferred by the individual mutations or the mutations conferred MICs
165 beyond the therapeutic window.

166 **WGS and single nucleotide polymorphism (SNP) calling**

167 WGS and data analysis was performed as previously described (29) and summarised in the
168 supplementary materials. The performance of WGS-based DST greatly depends on the
169 availability of robust markers of resistance. We therefore focussed on a set of high-
170 confidence resistance-associated genes (3, 14, 28) (Table 2). We additionally assessed the
171 impact of *eis* promotor mutations on amikacin and capreomycin resistance, as the association
172 of mutations in the *eis* promotor with resistance to the aforementioned drugs has been
173 reported but is not well established (11, 30).

174 **Ethics**

175 Local institutional review board or ethics committee approval was obtained at all local study
176 sites. Informed consent was obtained where requested per local regulations. This project was
177 also approved by the Cantonal Ethics Committee in Bern, Switzerland.

178 **Results**

179 Agreement between MGIT 960 and 7H10 agar dilution phenotypic DST

180 Table 3 and Figure 1 summarize the agreement between the semi-quantitative/quantitative
181 MIC determination by MGIT 960 and 7H10 agar dilution in terms of classifying strains as
182 belonging to the resistotype or wt populations as inferred by growth/no growth at the ECOFF
183 (Table 1). Agreement was high for all drugs, except ethambutol (see below). For most drugs,
184 the MGIT 960-based MICs were higher than the 7H10 agar dilution-based MICs. MICs
185 obtained using the two methods were within 1-2 two-fold dilution steps of each other. The
186 classifications into resistotype/wt populations demonstrated high rank correlations for most
187 drugs (Table 3 and Figure 1), except for capreomycin (supplementary Figure S4) due to few
188 strains demonstrating increased capreomycin MICs included in the study.

189 WGS and *in silico* resistance profile prediction

190 A total of 176 whole genome sequences with a median coverage of 676x (interquartile range
191 [IQR] = 37.48) were obtained. Median mapping percentage and percentage of genome
192 covered were 98.7 % (IQR = 0.94) and 99.4 % (IQR = 0.4), respectively. Genes involved in
193 drug resistance demonstrated high coverages with only 0.8 % of all positions suffering from
194 coverages below 7x (see supplementary materials). All major *M. tuberculosis* lineages,
195 except lineage 7, were represented in the study (L1 = 6, L2 = 36, L3 = 11, L4 = 123, L5 = 1,
196 L6 = 1). The strains showed a range of drug resistance profiles (Figure 2). Based on the set of
197 analysed genes (Table 2), 25 strains were predicted to be fully susceptible against all assayed
198 drugs, 59 strains were mono-/poly-resistant, 91 strains were MDR and two strains were
199 predicted to be extensively drug-resistant (XDR: isoniazid, rifampicin, fluoroquinolone and
200 aminoglycoside resistant).

201 Drug resistance profile prediction by WGS vs. phenotypic DST

202 After exclusion of known phylogenetic markers not involved in resistance, WGS-based
203 resistotype prediction using a defined set of target genes (Table 2) was highly congruent with
204 the categorical classification based on the phenotypic DST for most drugs (Table 3, Table 4,
205 Figure 1). Based on the *in silico* resistotype prediction, the MICs of mutant and wt strains
206 frequently followed a Gaussian distribution. Yet, the same resistance marker may confer
207 different MICs in different strains (supplementary Figures S1C, S2C, S3C, S8C, S9C S10C).
208 In some cases, the increase in the MIC conferred by a certain resistance mutation fell within

209 the distribution of the wt MIC (e.g. for *gidB*, *eis* promotor mutations, supplementary [Figures](#)
210 [S3C S6C](#)).

211 **Distinct wt and mutant MIC distributions**

212 MIC distributions of wt and mutant strains were well separated for rifampicin, rifabutin,
213 isoniazid, kanamycin A, amikacin, capreomycin, streptomycin and pyrazinamide, indicating
214 that the resistance markers used had a high positive predictive power (88.9 % overall positive
215 predictive power of associated with MIC increases). For streptomycin, two strains harboured
216 no mutations in the target genes, yet demonstrated high-level phenotypic resistance
217 (supplementary [Figure S3C](#)).

218 **Overlapping wt and mutant MIC distributions**

219 MIC distributions of wt and mutant strains overlapped for ethambutol, moxifloxacin and
220 ethionamide ([Figure 3](#)). For ethambutol and ethionamide, overlapping MIC distributions of
221 wt and mutant strains were associated with a large number of polymorphisms in resistance-
222 conferring genes (ethambutol resistance: 22 polymorphisms in *embB*, ethionamide resistance:
223 28 in *ethA*, 3 in *inhA*, 6 in *inhA* promoter). Solubility issues with ethionamide led to
224 quantitative differences in MGIT 960 vs. 7H10 agar dilution-based DST ([Table 3](#), [Figure 1](#)).
225 The overlap in MIC distributions between wt and strains carrying an *embB* mutation was
226 reduced by adjusting the critical concentration for ethambutol resistance from 5 mg/L to 2.5
227 mg/L (MGIT 960). However, there was variability in the MICs for the same mutation (e.g.
228 MIC *EmbB* M306I/V in 7H10 agar dilution: 4-16 mg/L –supplementary [Figure S2C](#)).
229 Moxifloxacin resistance was rare (n = 9, MGIT 960, critical concentration 0.25 mg/L) and
230 MIC distributions of mutant strains partially overlapped with those of wt. Sensitivity of the
231 genome-based moxifloxacin resistance prediction was 80.0 % ([Table 4](#)).

232 **Defining cut-offs for high-/low-level MICs**

233 **Isoniazid**

234 Mutations in the promoter of *inhA* caused low-level MICs <1 mg/L (7H10 agar dilution),
235 compared to strains harbouring mutations in *katG* or combinations of *inhA* promoter and
236 *katG* mutations which demonstrated MIC levels ranging from >1 mg/L to >32 mg/L in 7H10
237 agar dilution (supplementary [Figure S8C](#)). Defining cut-offs for low- (≤ 1 mg/L for MGIT

960/7H10 agar dilution) and high-level (>1 mg/L MGIT 960/7H10 agar dilution) isoniazid MICs is warranted.

Rifampicin/Rifabutin

Most mutations in *rpoB* increased the MIC for rifamycins beyond the therapeutic window (peak serum concentration 10 mg/L (28, 31)). However, some rare *rpoB* mutations (e.g. RpoB L452P, H445L – supplementary [Figure 9C](#)) demonstrated MICs within the therapeutic window. Defining cut-offs for low- and high-level MICs may thus be justified.

For rifampicin, cut-offs for low-/high-level MICs were $\leq 4/2$ mg/L for MGIT 960/7H10 agar dilution and $>4/2$ mg/L for MGIT 960/7H10 agar dilution, respectively.

For rifabutin, our data suggests a cut-off for low- and high-level MICs of $\leq 0.4/0.25$ or 0.5 mg/L for MGIT 960/7H10 agar dilution and $>0.4/0.25$ or 0.5 mg/L for MGIT 960/7H10 agar dilution, respectively.

Mutations in *rpoB* conferring resistance to rifampicin and rifabutin showed highly correlated increases ([Figure 4](#)) of MICs beyond the therapeutic window for most *rpoB* mutations ([Figure 3](#) and supplementary [Figure S9C](#) & [S10C](#)), indicating that both drugs are rendered clinically ineffective by the mutations identified in the dataset (32) and cannot substitute each other.

Amikacin

Few strains had mutations in the regions of *rrs* relevant for amikacin resistance or the *eis* promoter (n=12). Mutations in *rrs* were associated with high-level (>128 mg/L in 7H10 agar dilution) MICs. With regards to the *eis* promoter, only the C-14T mutation increased the MIC and led to low-level (2-4 mg/L in 7H10 agar dilution) MICs. The definition of a cut-off for low- (≤ 4 mg/L for MGIT 960/7H10 agar dilution) and high-level (4 mg/L for MGIT 960/7H10 agar dilution) amikacin MICs may be warranted.

Streptomycin

Certain mutations lead to MICs well beyond the therapeutic window (28) of streptomycin (e.g. RpsL K43R, MIC 7H10 agar dilution >128 mg/L, supplementary [Figure S3C](#)). On the other hand, *gidB* mutations increase the MIC only moderately (MIC 7H10 agar dilution 1-4 mg/L, supplementary [Figure 3C](#)). Mutational combinations in *gidB*, *rrs*, *rpsL* were common and produced a range of different MICs. Despite the distribution of MICs conferred by

267 combinations of mutations, there were distinct mutations that systematically lead to MICs
268 beyond the therapeutic window, e.g. RpsL K43R. Defining a cut-off for low-level (MGIT
269 960 ≤ 4 mg/L, 7H10 agar dilution ≤ 4 -8 mg/L) and high-level streptomycin MICs (MGIT 960
270 > 4 mg/L, 7H10 agar dilution > 4 -8 mg/L) is warranted.

271 Discussion

272 The results of MGIT 960 and 7H10 agar dilution-based phenotypic DST methods were
273 highly correlated and suitable to separate the resistotype from the wt populations. Based on
274 phenotypic DST results and WGS, we were able to define cut-offs for high- and low-level
275 MICs for isoniazid, rifampicin, streptomycin and amikacin. Defining such cut-offs may serve
276 as starting points for correlating mutational, DST and pharmaco-kinetic/dynamic data to gain
277 more insight into the influence of individual mutations on treatment outcomes, especially in
278 the light of e.g. increased drug dosing.

279 Our data suggest that the current WHO-defined critical concentration for phenotypic DST of
280 ethambutol by MGIT 960 (5 mg/L) is too high and may misclassify strains as belonging to
281 the wt population when compared to the 7H10 agar dilution-based classification. Given the
282 narrow therapeutic window for ethambutol, this may lead to mistreatment due to presumed
283 ethambutol susceptibility. After adjusting the ECOFF to 2.5 mg/L for MGIT 960, we
284 observed a strong improvement of the categorical agreement between MGIT 960- and 7H10
285 agar dilution-based classification into resistotype/wt populations.

286 The mutations identified by WGS had a high predictive power to classify strains as belonging
287 to the resistotype population. However, the predictive power depends on a number of factors.
288 For instance, the increase in MIC conferred by an identical mutation can vary greatly in
289 different strains (e.g. EmbB M306I/V, RpsL K88R) (33). Such variation may be clinically
290 relevant if there is a significant overlap between the MICs of mutant and wt strains (23), as
291 was the case for strains harbouring mutations in genes associated with ethionamide,
292 ethambutol and streptomycin (e.g. *gidB*) resistance. . Furthermore, it is difficult to classify
293 strains as part of resistotype or wt populations if the MIC increase lies within the therapeutic
294 window of a drug. The overlap between MICs of mutant and wt strains is confounded by the
295 fact that we only screened for mutations in genes which had previously been associated with
296 drug resistance. We might thus have missed possible resistance-conferring mutations in other
297 genes. Additionally, WGS will always produce distributions of coverages, which in term will

298 inevitably lead to certain regions in the genome suffering from low coverage, preventing the
299 detection of mutations. However, in cases where we observed elevated MICs without any
300 mutations detected the target genes, coverage issues could not explain the absence of any
301 mutations. Furthermore, the strain genetic background (34), non-mutational mechanisms (e.g.
302 modulation of gene expression) (35), as well as drug efflux mechanisms (36) may contribute
303 to the variability in increase of the MIC conferred by resistance mutations.

304 The predictive power of mutations in target genes also depends on removing phylogenetic
305 markers not involved in increasing MICs. Separating phylogenetic from resistance-associated
306 markers works well for essential (highly conserved) genes such as *rpoB*, *rpsL*, *rrs* but is
307 problematic in non-essential genes involved in the conversion of prodrugs into their active
308 forms like *pncA* (pyrazinamide), *ethA* (ethionamide) or in genes that generally exhibit higher
309 numbers of polymorphisms e.g. *embB*. Of note, the *embABC* operon is highly polymorphic,
310 harbouring more polymorphisms than expected by chance (mutations in *embABC* operon =
311 81, expected = 44.8, $p = 9.174e-07$, binomial test). Mutations conferring increased ethambutol
312 MICs (37) will therefore inevitably evolve in the presence of phylogenetic SNPs and may
313 interact epistatically to produce the variability in MICs we observed for wt strains and for the
314 most common marker associated with increased ethambutol resistance MICs, *embB*
315 M306I/V. The *embABC* operon is involved in the biosynthesis of decaprenylphosphoryl- β -d-
316 arabinose, which is an integral component of the mycobacterial cell wall. The cell envelope
317 interacts with the host immune system and the high levels of diversity of these genes might
318 be the product of diversifying selection due to host immune pressure. The influence of
319 polymorphisms in the *embABC* operon on MICs in general is supported by the observation
320 that sub-inhibitory concentrations of ethambutol lower the MICs for isoniazid, rifampicin and
321 streptomycin (38). Even small changes in activity of the decaprenylphosphoryl- β -d-arabinose
322 biosynthetic and utilisation pathway might thus alter cell wall permeability and influence
323 MICs of several drugs.

324 Similarly, in the case of increased streptomycin MICs, the RpsL substitution K88R exhibited
325 a wide range of MIC increases, partially within the therapeutic window of the drug.
326 Streptomycin was the first effective antituberculous drug discovered (39) and has been used
327 for decades. The long-term use has produced complex resistance profiles with multiple
328 mutations known to increase streptomycin MICs on their own (e.g. in *gidB*, *rpsL*, *rrs*)
329 occurring concomitantly, producing wide ranges of MICs. Furthermore, many strains with
330 increased streptomycin MICs displayed MDR/XDR phenotypes. Mutations conferring

331 increased streptomycin MICs are frequently found in backgrounds which have mutations in
332 genes affecting the information pathway (DNA → RNA → proteins) – e.g. *gyrA* (DNA
333 gyrase), *rpoB* (DNA-dependent RNA polymerase), *rrs* (ribosomal RNA). The simultaneous
334 presence of multiple MIC increasing mutations may alter the adaptive landscape (40, 41). In
335 addition, non-mutational processes (e.g. alteration of gene expression) may compensate for
336 fitness costs of drug resistance and at the same time alter the MIC for the drug (35). This has
337 not been demonstrated for streptomycin resistance in *M. tuberculosis*, but it seems possible
338 that compensation of fitness costs in MDR phenotypes might alter the MIC for streptomycin
339 (40), considering that streptomycin is not part of the current standard treatment regimen and
340 selection for high-level streptomycin MICs is relaxed.

341 Concerning *eis* promoter mutations and aminoglycoside resistance, there is mounting
342 evidence that the *eis* C-14T promotor mutation may confer clinically relevant increases in
343 amikacin MICs, especially in the light of the revised critical concentrations for amikacin (2
344 mg/L, 7H10 agar dilution) [11].

345 We observed an overrepresentation of lineage 4 and 2 strains in our sample set. The strain set
346 used to establish the methodology was collated with a specific aim to include drug-resistant
347 strains and given the frequent association of lineage 2 and 4 with drug-resistance (42, 43), the
348 observed skew is not surprising. Furthermore, lineage 2 and 4 strains are also frequently
349 isolated at the collection sites of the strain set used to apply the methodology (Ivory Coast,
350 Peru and Democratic Republic of the Congo). Similarly, increased MICs for a number of
351 drugs (amikacin, capreomycin, kanamycin, moxifloxacin) was rare, reflecting the scarcity of
352 pre-XDR/XDR phenotypes in Switzerland and at the sites of prospective sampling.

353 With 63.6 % – 80.8 %, sensitivities were low (44) for a number of drugs (i.e. for amikacin,
354 moxifloxacin, pyrazinamide) (table 4), but were comparable to other studies not employing a
355 database of pre-defined resistance mutations (14, 17, 45). The observed low sensitivities for
356 some drugs were either due to few strains belonging to the resistotype population included in
357 the dataset, the presence of additional resistance mutations in genes not assessed or due to
358 unknown resistance mechanisms and not due to low coverages prohibiting the detection of
359 mutations. The use of a curated SNP-database containing high-confidence drug-resistance
360 mutations would improve sensitivity for some drugs where additional targets, less well
361 associated with MIC increases, are known (44, 46). However, reliance on a predefined
362 resistance mutation database comes at the cost of reduced sensitivity. After known

363 phylogenetic mutations have been removed, it is important to treat any mutation in known
364 target genes as potentially involved in drug resistance. In cases where previously unknown
365 mutations (i.e. neither known to increase MICs, nor a known phylogenetic SNP) in
366 resistance-related genes are detected, genetic engineering and targeted DST is necessary to
367 confirm or reject the drug-resistance conferring nature of a novel mutation to achieve high
368 sensitivities and specificities for whole genome sequencing-based DST.

369 Generating high-quality quantitative DST data using diverse *M. tuberculosis* strains is
370 important to accurately define the ECOFF and subsequently guide treatment decisions. The
371 two quantitative DST methods employed are difficult to standardize across laboratories,
372 technically demanding and at best challenging to scale up. Microtiter plate-based quantitative
373 DST methods (47, 48) have the potential to aid in the generation of more high quality DST
374 data due to their standardized formulation and relative ease of application compared to
375 established methods.

376 In conclusion, we demonstrate that MGIT 960 and 7H10 agar dilution-based phenotypic DST
377 provide highly congruent classifications of strains into resistotype or wt populations. WGS
378 has high predictive power to infer resistance profiles without the need for time-consuming
379 phenotypic methods. Limitations due to overlapping distributions of wt and resistotype MICs,
380 varying MICs for the same mutations in different strains, presence of phylogenetic markers in
381 resistance-associated genes and rare resistance markers with low frequencies will likely be
382 resolved by on-going large-scale projects (e.g. ReSeqTB and others (15, 49)) combining
383 phenotypic DST with WGS of thousands of *M. tuberculosis* isolates. Our findings, together
384 with those of on-going studies will pave the way for the replacement of phenotypic DST with
385 drug resistance profile prediction based on WGS in the coming years.

386 **Acknowledgements**

387 We would like to thank Alexandra Mushegian for critically reading the manuscript and
388 improving the writing. Whole genome analysis was performed at sciCORE
389 (<http://scicore.unibas.ch/>) scientific computing core facility at University of Basel.

390 **Funding**

391 This work was supported by the European Research Council [grant number 309540-
392 EVODRTB], the Swiss National Science Foundation [IZRJZ3_164171, 310030-166687,

393 IZLSZ3_170834, CRSII5_177163, 31003A_153349 and 320030_153442/1, special project
394 funding 174281] and SystemsX.ch. The International epidemiology Databases to Evaluate
395 AIDS (IeDEA) is supported by the National Institutes of Allergy and Infectious Diseases
396 (NIAID) under award numbers U01 AI096299, U01 AI069919, U01 AI069924, U01
397 AI069911, U01 AI069907, U01 AI096186, and U01 AI069923. The content of this
398 publication is solely the responsibility of the authors and does not necessarily represent the
399 official views of any of the funding agencies mentioned above.

400

401 **Conflict of interest**

402 Peter M. Keller reports travel grants by Copan Italia SpA outside of the submitted work. Erik
403 C. Böttger is a consultant for AID Diagnostics.

404

405 **Tables**

406

407

408

409

410

411

412

413

414

415

416

417

Table 1 Epidemiological cut-offs (ECOFF) used for 7H10 agar dilution and MGIT 960 phenotypic DST, derived from wild-type MIC distributions determined in this study. The values given in parentheses are the critical concentrations recommended by the WHO in 2014 [22].

Antibiotic	ECOFF agar dilution (mg/L)	ECOFF MGIT 960 (mg/L)
Ethionamide	1 (5)	5
Ethambutol	2 (5)	5
Capreomycin	4	2.5
Streptomycin	0.5 (2)	1
Kanamycin A	2 (5)	2 (2.5)
Amikacin	1 (4)	1
Moxifloxacin	0.25 (0.5)	0.25 (0.5)
Isoniazid	0.125 (0.2)	0.1
Rifampicin	0.5 (1)	1
Rifabutin	0.0625	0.1
Pyrazinamide	NA	100

Table 2 List of genes implicated in drug resistance in *M. tuberculosis* which were screened for polymorphisms by WGS. List adapted from [3, 12, 23].

Drug	Target gene(s)
Ethionamide	<i>ethA</i> , <i>inhA</i> , <i>inhA</i> promoter
Ethambutol	<i>embB</i>
Capreomycin	<i>rrs</i> , <i>eis</i> promoter, <i>tlyA</i>
Streptomycin	<i>rrs</i> , <i>gidB</i> , <i>rpsL</i>
Kanamycin A	<i>rrs</i> , <i>eis</i> promoter
Amikacin	<i>rrs</i> , <i>eis</i> promoter
Moxifloxacin	<i>gyrA</i>
Isoniazid	<i>katG</i> , <i>inhA</i> promoter
Rifampicin/rifabutin	<i>rpoB</i>
Pyrazinamide	<i>pncA</i> , <i>pncA</i> promoter

Table 3 Summary statistics of the method agreement between 7H10 agar dilution- and MGIT 960-based phenotypic DST for all drugs assayed in this study.

Antibiotic	n	Categorical agreement (%)	SD of $\log_2(\text{MIC MGIT 960/MIC agar dilution})$	γ
Ethionamide	56	95	1.9 ± 0.3	0.91
Ethambutol	171	73	1.9 ± 0.5	0.94
Capreomycin	56	98	1.5 ± 0.5	0.65
Streptomycin	56	93	1.5 ± 0.3	0.98
Kanamycin A	56	98	1.2 ± 0.2	0.8
Amikacin	174	98	1.4 ± 0.6	1
Moxifloxacin	173	99	1 ± 0.2	1
Isoniazid	173	96	1.2 ± 0.1	1
Rifampicin	174	99	NA	1
Rifabutin	56	96	0.8 ± 0.1	0.98

Table 4 Sensitivity and specificity of the genome-based drug resistance profile prediction using the 7H10 agar dilution-based categorical classification as the gold standard for all drugs except pyrazinamide, for which the MGIT 960 categorical classification was used.

Drug	Sensitivity (%)	Specificity (%)
Ethionamide	75.0	92.9
Ethambutol	89.6	94.2
Capreomycin	75.0	94
Streptomycin	68.0	92.1
Kanamycin A	83.3	98.8
Amikacin	63.6	96.9
Moxifloxacin	80.0	90.2
Isoniazid	93.6	96.8
Rifampicin	100	94.0
Rifabutin	98.9	94.0
Pyrazinamide	80.8	88.9

Figure legends

Figure 1

Method agreement between phenotypic DST performed with MGIT 960 and 7H10 agar dilution represented as Bland-Altman plots for all drugs tested in this study.

Figure 2

Maximum likelihood phylogeny of 176 *M. tuberculosis* strains based on 20510 variable positions. Reference strains labeled with green tip labels. Main lineages are highlighted as follows: red L4, purple L3, blue L2, pink L1, green L6, brown L5. Scale bar indicates number of substitutions per site. Phylogeny rooted on *M. canettii*. Colored bars indicate resistance

445 mutations per gene and within a distinct column (gene) each colored bar represents a distinct
446 mutation. Black bars indicate no mutation, i.e. wt.

447

448 **Figure 3**

449 Histograms of MICs (7H10 agar dilution) for all drugs assayed in this study

450

451 **Figure 4**

452 Correlation between 7H10 agar dilution MICs for rifampicin and rifabutin

References

1. World Health Organization. 2010. Treatment of tuberculosis: Guidelines Treatment of Tuberculosis: Guidelines 4th Editio.
2. Domínguez J, Boettger EC, Cirillo D, Cobelens F, Eisenach KD, Gagneux S, Hillemann D, Horsburgh R, Molina-Moya B, Niemann S, Tortoli E, Whitelaw A, Lange C, TBNET, RESIST-TB networks. 2016. Clinical implications of molecular drug resistance testing for Mycobacterium tuberculosis: a TBNET/RESIST-TB consensus statement. *Int J Tuberc Lung Dis* 20:24–42.
3. Gygli SM, Borrell S, Trauner A, Gagneux S. 2017. Antimicrobial resistance in Mycobacterium tuberculosis: mechanistic and evolutionary perspectives. *FEMS Microbiol Rev* 41:354–373.
4. Deggin-Messmer V, Bloemberg G V., Ritter C, Voit A, Hömke R, Keller PM, Böttger EC. 2016. Diagnostic Molecular Mycobacteriology in Regions With Low Tuberculosis Endemicity: Combining Real-time PCR Assays for Detection of Multiple Mycobacterial Pathogens With Line Probe Assays for Identification of Resistance Mutations. *EBioMedicine* 9:228–237.
5. Nathavitharana RR, Hillemann D, Schumacher SG, Schlueter B, Ismail N, Omar SV, Sikhondze W, Havumaki J, Valli E, Boehme C, Denking CM. 2016. Multicenter Noninferiority Evaluation of Hain GenoType MTBDR plus Version 2 and Nipro NTM+MDRTB Line Probe Assays for Detection of Rifampin and Isoniazid Resistance. *J Clin Microbiol* 54:1624–1630.
6. Ritter C, Lucke K, Sirgel FA, Warren RW, van Helden PD, Böttger EC, Bloemberg G V. 2014. Evaluation of the AID TB Resistance Line Probe Assay for Rapid Detection of Genetic Alterations Associated with Drug Resistance in Mycobacterium tuberculosis Strains. *J Clin Microbiol* 52:940–946.
7. World Health Organization. 2016. WHO Guideline: The use of molecular line probe assays for the detection of resistance to isoniazid and rifampicin. Geneva.
8. WHO. 2013. Automated Real-Time Nucleic Acid Amplification Technology for Rapid and Simultaneous Detection of Tuberculosis and Rifampicin Resistance: Xpert MTB/RIF Assay for the Diagnosis of Pulmonary and Extrapulmonary TB in Adults and Children: Policy Update Automated Real-Time Nucleic Acid Amplification Technology for Rapid and Simultaneous Detection of Tuberculosis and Rifampicin Resistance: Xpert MTB/RIF Assay for the Diagnosis of Pulmonary and Extrapulmonary TB in Adults and Children: Policy Update.
9. Engström A. 2016. Fighting an old disease with modern tools: characteristics and molecular detection methods of drug-resistant Mycobacterium tuberculosis. *Infect Dis (Auckl)* 48:1–17.
10. Streicher EM, Bergval I, Dheda K, Böttger EC, Gey van Pittius NC, Bosman M, Coetzee G, Anthony RM, van Helden PD, Victor TC, Warren RM. 2012. Mycobacterium tuberculosis Population Structure Determines the Outcome of Genetics-Based Second-Line Drug Resistance Testing. *Antimicrob Agents Chemother* 56:2420–2427.

- 495 11. World Health Organization. 2018. Technical Report on critical concentrations for drug
496 susceptibility testing of medicines used in the treatment of drug-resistant tuberculosis.
- 497 12. World Health Organization. 2018. Technical manual for drug susceptibility testing of
498 medicines used in the treatment of tuberculosis. Geneva.
- 499 13. Coll F, McNerney R, Preston MD, Guerra-Assunção JA, Warry A, Hill-Cawthorne G,
500 Mallard K, Nair M, Miranda A, Alves A, Perdigão J, Viveiros M, Portugal I, Hasan Z,
501 Hasan R, Glynn JR, Martin N, Pain A, Clark TG. 2015. Rapid determination of anti-
502 tuberculosis drug resistance from whole-genome sequences. *Genome Med* 7:51.
- 503 14. Walker TM, Kohl TA, Omar S V, Hedge J, Del Ojo Elias C, Bradley P, Iqbal Z,
504 Feuerriegel S, Niehaus KE, Wilson DJ, Clifton DA, Kapatai G, Ip CLC, Bowden R,
505 Drobniewski FA, Allix-Béguec C, Gaudin C, Parkhill J, Diel R, Supply P, Crook DW,
506 Smith EG, Walker AS, Ismail N, Niemann S, Peto TEA. 2015. Whole-genome
507 sequencing for prediction of *Mycobacterium tuberculosis* drug susceptibility and
508 resistance: a retrospective cohort study. *Lancet Infect Dis* 15:1193–1202.
- 509 15. The CRyPTIC Consortium and the 100000 Genomes Project. 2018. Prediction of
510 Susceptibility to First-Line Tuberculosis Drugs by DNA Sequencing. *N Engl J Med*
511 *NEJMoa1800474*.
- 512 16. Dookie N, Rambaran S, Padayatchi N, Mahomed S, Naidoo K. 2018. Evolution of
513 drug resistance in *Mycobacterium tuberculosis*: a review on the molecular
514 determinants of resistance and implications for personalized care. *J Antimicrob*
515 *Chemother*.
- 516 17. Shea J, Halse TA, Lapierre P, Shudt M, Kohlerschmidt D, Van Roey P, Limberger R,
517 Taylor J, Escuyer V, Musser KA. 2017. Comprehensive Whole-Genome Sequencing
518 and Reporting of Drug Resistance Profiles on Clinical Cases of *Mycobacterium*
519 *tuberculosis* in New York State. *J Clin Microbiol* 55:1871–1882.
- 520 18. Colman RE, Anderson J, Lemmer D, Lehmkühl E, Georghiou SB, Heaton H, Wiggins
521 K, Gillette JD, Schupp JM, Catanzaro DG, Crudu V, Cohen T, Rodwell TC,
522 Engelthaler DM. 2016. Rapid Drug Susceptibility Testing of Drug Resistant
523 *Mycobacterium tuberculosis* Directly from Clinical Samples using Amplicon
524 Sequencing: A Proof of Concept Study. *J Clin Microbiol* 54:JCM.00535-16.
- 525 19. Springer B, Lucke K, Calligaris-Maibach R, Ritter C, Bottger EC. 2009. Quantitative
526 Drug Susceptibility Testing of *Mycobacterium tuberculosis* by Use of MGIT 960 and
527 EpiCenter Instrumentation. *J Clin Microbiol* 47:1773–1780.
- 528 20. Stucki D, Ballif M, Egger M, Furrer H, Altpeter E, Battegay M, Droz S, Bruderer T,
529 Coscolla M, Borrell S, Zürcher K, Janssens J-P, Calmy A, Mazza Stalder J, Jaton K,
530 Rieder HL, Pfyffer GE, Siegrist HH, Hoffmann M, Fehr J, Dolina M, Frei R,
531 Schrenzel J, Böttger EC, Gagneux S, Fenner L. 2016. Standard Genotyping
532 Overestimates Transmission of *Mycobacterium tuberculosis* among Immigrants in a
533 Low-Incidence Country. *J Clin Microbiol* 54:1862–1870.
- 534 21. Bloemberg G V., Keller PM, Stucki D, Trauner A, Borrell S, Latshang T, Coscolla M,
535 Rothe T, Hömke R, Ritter C, Feldmann J, Schulthess B, Gagneux S, Böttger EC. 2015.
536 Acquired Resistance to Bedaquiline and Delamanid in Therapy for Tuberculosis. *N*
537 *Engl J Med* 373:1986–1988.

- 538 22. Egger M, Ekouevi DK, Williams C, Lyamuya RE, Mukumbi H, Braitstein P, Hartwell
539 T, Graber C, Chi BH, Boule A, Dabis F, Wools-Kaloustian K. 2012. Cohort Profile:
540 The international epidemiological databases to evaluate AIDS (IeDEA) in sub-Saharan
541 Africa. *Int J Epidemiol* 41:1256–1264.
- 542 23. Schön T, Miotto P, Köser CU, Viveiros M, Böttger E, Cambau E. 2017.
543 *Mycobacterium tuberculosis* drug-resistance testing: challenges, recent developments
544 and perspectives. *Clin Microbiol Infect* 23:154–160.
- 545 24. Ängeby K, Juréen P, Kahlmeter G, Hoffner S, Schön T. 2012. Challenging a dogma:
546 antimicrobial susceptibility testing breakpoints for *Mycobacterium tuberculosis*. *Bull*
547 *World Health Organ* 90:693–698.
- 548 25. World Health Organization. 2014. Companion handbook to the WHO guidelines for
549 the programmatic management of drug-resistant tuberculosis. Geneva.
- 550 26. Delignette-Muller ML, Dutang C. 2015. *fitdistrplus*: an R package for fitting
551 distributions. *J Stat Softw* 64:1–34.
- 552 27. Martin Bland J, Altman D. 1986. STATISTICAL METHODS FOR ASSESSING
553 AGREEMENT BETWEEN TWO METHODS OF CLINICAL MEASUREMENT. *Lancet*
554 327:307–310.
- 555 28. Böttger EC. 2011. The ins and outs of *Mycobacterium tuberculosis* drug susceptibility
556 testing. *Clin Microbiol Infect* 17:1128–1134.
- 557 29. Ghielmetti G, Coscolla M, Ruetten M, Friedel U, Loiseau C, Feldmann J, Steinmetz
558 HW, Stucki D, Gagneux S. 2017. Tuberculosis in Swiss captive Asian elephants:
559 microevolution of *Mycobacterium tuberculosis* characterized by multilocus variable-
560 number tandem-repeat analysis and whole-genome sequencing. *Sci Rep* 7:14647.
- 561 30. Kambli P, Ajbani K, Nikam C, Sadani M, Shetty A, Udawadia Z, Georghiou SB,
562 Rodwell TC, Catanzaro A, Rodrigues C. 2016. Correlating *rrs* and *eis* promoter
563 mutations in clinical isolates of *Mycobacterium tuberculosis* with phenotypic
564 susceptibility levels to the second-line injectables. *Int J Mycobacteriology* 5:1–6.
- 565 31. Sekaggya-Wiltshire C, von Braun A, Lamorde M, Ledergerber B, Buzibye A, Henning
566 L, Musaazi J, Gutteck U, Denti P, de Kock M, Jetter A, Byakika-Kibwika P, Eberhard
567 N, Matovu J, Joloba M, Muller D, Manabe YC, Kamya MR, Corti N, Kambugu A,
568 Castelnovo B, Fehr JS. 2018. Delayed Sputum Culture Conversion in Tuberculosis–
569 Human Immunodeficiency Virus–Coinfected Patients With Low Isoniazid and
570 Rifampicin Concentrations. *Clin Infect Dis* 67:708–716.
- 571 32. Berrada ZL, Lin S-YG, Rodwell TC, Nguyen D, Schecter GF, Pham L, Janda JM,
572 Elmarachli W, Catanzaro A, Desmond E. 2016. Rifabutin and rifampin resistance
573 levels and associated *rpoB* mutations in clinical isolates of *Mycobacterium*
574 *tuberculosis* complex. *Diagn Microbiol Infect Dis* 85:177–181.
- 575 33. Ruesen C, Riza AL, Florescu A, Chaidir L, Editoiu C, Aalders N, Nicolosu D, Grecu
576 V, Ioana M, van Crevel R, van Ingen J. 2018. Linking minimum inhibitory
577 concentrations to whole genome sequence-predicted drug resistance in *Mycobacterium*
578 *tuberculosis* strains from Romania. *Sci Rep* 8:9676.

- 579 34. Fenner L, Egger M, Bodmer T, Altpeter E, Zwahlen M, Jaton K, Pfyffer GE, Borrell S,
580 Dubuis O, Bruderer T, Siegrist HH, Furrer H, Calmy A, Fehr J, Stalder JM, Ninet B,
581 Böttger EC, Gagneux S. 2012. Effect of Mutation and Genetic Background on Drug
582 Resistance in *Mycobacterium tuberculosis*. *Antimicrob Agents Chemother* 56:3047–
583 3053.
- 584 35. Freihofer P, Akbergenov R, Teo Y, Juskeviciene R, Andersson DI, Böttger EC. 2016.
585 Nonmutational compensation of the fitness cost of antibiotic resistance in
586 mycobacteria by overexpression of tlyA rRNA methylase. *RNA* 22:1836–1843.
- 587 36. da Silva PEA, Machado D, Ramos D, Couto I, Von Groll A, Viveiros M. 2016. Efflux
588 Pumps in *Mycobacteria*: Antimicrobial Resistance, Physiological Functions, and Role
589 in Pathogenicity, p. 527–559. *In* Li, X-Z, Elkins, CA, Zgurskaya, HI (eds.), *Efflux-*
590 *Mediated Antimicrobial Resistance in Bacteria*. Springer International Publishing,
591 Cham.
- 592 37. Safi H, Lingaraju S, Amin A, Kim S, Jones M, Holmes M, McNeil M, Peterson SN,
593 Chatterjee D, Fleischmann R, Alland D. 2013. Evolution of high-level ethambutol-
594 resistant tuberculosis through interacting mutations in decaprenylphosphoryl- β -D-
595 arabinose biosynthetic and utilization pathway genes. *Nat Genet* 45:1190–1197.
- 596 38. Jagannath C, Reddy VM, Gangadharam PRJ. 1995. Enhancement of drug
597 susceptibility of multi-drug resistant strains of *Mycobacterium tuberculosis* by
598 ethambutol and dimethyl sulphoxide. *J Antimicrob Chemother* 35:381–390.
- 599 39. Schatz A, Bugle E, Waksman SA. 1944. Streptomycin, a Substance Exhibiting
600 Antibiotic Activity Against Gram-Positive and Gram-Negative Bacteria. *Exp Biol Med*
601 55:66–69.
- 602 40. Moura de Sousa J, Balbontín R, Durão P, Gordo I. 2017. Multidrug-resistant bacteria
603 compensate for the epistasis between resistances. *PLOS Biol* 15:e2001741.
- 604 41. Borrell S, Teo Y, Giardina F, Streicher EM, Kloppe M, Feldmann J, Müller B, Victor
605 TC, Gagneux S. 2013. Epistasis between antibiotic resistance mutations drives the
606 evolution of extensively drug-resistant tuberculosis. *Evol Med Public Heal* 2013:65–
607 74.
- 608 42. Casali N, Nikolayevskyy V, Balabanova Y, Harris SR, Ignatyeva O, Kontsevaya I,
609 Corander J, Bryant J, Parkhill J, Nejentsev S, Horstmann RD, Brown T, Drobniewski
610 F. 2014. Evolution and transmission of drug-resistant tuberculosis in a Russian
611 population. *Nat Genet* 46:279–86.
- 612 43. Cohen KA, Abeel T, Manson McGuire A, Desjardins CA, Munsamy V, Shea TP,
613 Walker BJ, Bantubani N, Almeida D V., Alvarado L, Chapman SB, Mvelase NR,
614 Duffy EY, Fitzgerald MG, Govender P, Gujja S, Hamilton S, Howarth C, Larimer JD,
615 Maharaj K, Pearson MD, Priest ME, Zeng Q, Padayatchi N, Grosset J, Young SK,
616 Wortman J, Mlisana KP, O'Donnell MR, Birren BW, Bishai WR, Pym AS, Earl AM.
617 2015. Evolution of Extensively Drug-Resistant Tuberculosis over Four Decades:
618 Whole Genome Sequencing and Dating Analysis of *Mycobacterium tuberculosis*
619 Isolates from KwaZulu-Natal. *PLOS Med* 12:e1001880.
- 620 44. World Health Organization. 2018. The use of next-generation sequencing technologies
621 for the detection of mutations associated with drug resistance in *Mycobacterium*

- 622 tuberculosis complex: technical guide. Geneva.
- 623 45. Farhat MR, Sultana R, Iartchouk O, Bozeman S, Galagan J, Sisk P, Stolte C,
624 Nebenzahl-Guimaraes H, Jacobson K, Sloutsky A, Kaur D, Posey J, Kreiswirth BN,
625 Kurepina N, Rigouts L, Streicher EM, Victor TC, Warren RM, van Soolingen D,
626 Murray M. 2016. Genetic Determinants of Drug Resistance in *Mycobacterium*
627 tuberculosis and Their Diagnostic Value. *Am J Respir Crit Care Med* 194:621–30.
- 628 46. Miotto P, Tessema B, Tagliani E, Chindelevitch L, Starks AM, Emerson C, Hanna D,
629 Kim PS, Liwski R, Zignol M, Gilpin C, Niemann S, Denking CM, Fleming J,
630 Warren RM, Crook D, Posey J, Gagneux S, Hoffner S, Rodrigues C, Comas I,
631 Engelthaler DM, Murray M, Alland D, Rigouts L, Lange C, Dheda K, Hasan R,
632 Ranganathan UDK, McNerney R, Ezewudo M, Cirillo DM, Schito M, Köser CU,
633 Rodwell TC. 2017. A standardised method for interpreting the association between
634 mutations and phenotypic drug resistance in *Mycobacterium tuberculosis*. *Eur Respir J*
635 50:1701354.
- 636 47. Lee J, Armstrong DT, Ssengooba W, Park J-A, Yu Y, Mumbowa F, Namaganda C,
637 Mboowa G, Nakayita G, Armakovitch S, Chien G, Cho S-N, Via LE, Barry CE, Ellner
638 JJ, Alland D, Dorman SE, Joloba ML. 2014. Sensititre MYCOTB MIC Plate for
639 Testing *Mycobacterium tuberculosis* Susceptibility to First- and Second-Line Drugs.
640 *Antimicrob Agents Chemother* 58:11–18.
- 641 48. Rancoita PM V., Cugnata F, Gibertoni Cruz AL, Borroni E, Hoosdally SJ, Walker TM,
642 Grazian C, Davies TJ, Peto TEA, Crook DW, Fowler PW, Cirillo DM, Crook DW,
643 Peto TEA, Walker AS, Hoosdally SJ, Gibertoni Cruz AL, Grazian C, Walker TM,
644 Fowler PW, Wilson D, Clifton D, Iqbal Z, Hunt M, Smith EG, Rathod P, Jarrett L,
645 Matias D, Cirillo DM, Borroni E, Battaglia S, Chiacchiaretta M, De Filippo M,
646 Cabibbe A, Tahseen S, Mistry N, Nilgiriwala K, Chitalia V, Ganesan N, Papewar A,
647 Rodrigues C, Kambli P, Surve U, Khot R, Niemann S, Kohl T, Merker M, Hoffmann
648 H, Lehmann S, Plesnik S, Ismail N, Omar SV, Joseph L, Marubini E, Thwaites G,
649 Thuy Thuong TN, Ngoc NH, Srinivasan V, Moore D, Coronel J, Solano W, He G, Zhu
650 B, Zhou Y, Ma A, Yu P, Schito M, Claxton P, Laurenson I. 2018. Validating a 14-
651 Drug Microtiter Plate Containing Bedaquiline and Delamanid for Large-Scale
652 Research Susceptibility Testing of *Mycobacterium tuberculosis*. *Antimicrob Agents*
653 *Chemother* 62:1–15.
- 654 49. Starks AM, Avilés E, Cirillo DM, Denking CM, Dolinger DL, Emerson C, Gallarda
655 J, Hanna D, Kim PS, Liwski R, Miotto P, Schito M, Zignol M. 2015. Collaborative
656 Effort for a Centralized Worldwide Tuberculosis Relational Sequencing Data Platform.
657 *Clin Infect Dis* 61Suppl 3:S141-6.

658




Figure 1 Method agreement between phenotypic DST performed with MGIT 960 and 7H10 agar dilution represented as Bland-Altman plots for all drugs tested in this study.




Figure 2 Maximum likelihood phylogeny of 176 *M. tuberculosis* strains based on 20510 variable positions. Reference strains labeled with green tip labels. Main lineages are highlighted as follows: red L4, purple L3, blue L2, pink L1, green L6, brown L5. Scale bar indicates number of substitutions per site. Phylogeny rooted on *M. canettii*. Colored bars indicate resistance mutations per gene and within a distinct column (gene) each colored bar represents a distinct mutation. Black bars indicate no mutation, i.e. wt.




Figure 3 Histograms of MICs (7H10 agar dilution) for all drugs assayed in this study




Figure 4 Correlation between 7H10 agar dilution MICs for rifampicin and rifabutin